



LAWRENCE  
LIVERMORE  
NATIONAL  
LABORATORY

# The Utilization of Spin Polarized Photoelectron Spectroscopy as a Probe of Electron Correlation with an Ultimate Goal of Pu

J. G. Tobin, S. W. Yu, B. W. Chung, S. A. Morton,  
T. Komesu, G. D. Waddill

February 11, 2008

Symposium Proceedings of the Material Research Society

## **Disclaimer**

---

This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.

## The Utilization of Spin Polarized Photoelectron Spectroscopy as a Probe of Electron Correlation with an Ultimate Goal of Pu

J.G. Tobin<sup>1</sup>, S.W. Yu<sup>1</sup>, B.W. Chung<sup>1</sup>, S.A. Morton<sup>1,2</sup>,  
T. Komesu<sup>3,\*</sup> and G.D. Waddill<sup>3</sup>

1. Lawrence Livermore National Laboratory, LLNS-LLC, Livermore, CA, USA

2. Lawrence Berkeley National Laboratory, Berkeley, CA, USA

3. University of Missouri-Rolla, Department of Physics, Rolla, MO, USA

We are developing the technique of spin-polarized photoelectron spectroscopy as a probe of electron correlation with the ultimate goal of resolving the Pu electronic structure controversy. Over the last several years, we have demonstrated the utility of spin polarized photoelectron spectroscopy for determining the fine details of the electronic structure in complex systems such as those shown below.

The proposed half-metallic ferro-magnet Fe<sub>3</sub>O<sub>4</sub> [Figure 1, Ref 2]




Figure 1  
a) Spin resolved photoemission spectra from a) "as received" sample  $h\nu=160\text{eV}$ .  
b) Spin resolved photoemission spectra from a sample after Ne<sup>+</sup> sputtering,  $h\nu=160\text{eV}$ .  
Note the large effect observed in the as received sample in (a) and the reduction of the effect in (b)

The 2p core levels of ultra-thin films such as Fe [Figure 2, Ref 3].




Figure 2  
Simulated spin-resolved spectrum of the Fe 2p doublet, following Reference 3.  
Red is spin up and blue is spin down.

The valence states of Fe on compound semiconductors [Figure 3, Ref 4].




Figure 3

An example of some data collected on Beamline 7 at the ALS is shown here. The photon energy was 145 eV and the photon polarization is linear. The spin-resolved valence band photoemission results are from Fe overlayers, as part of a larger collaboration. [4] The Fermi edge is near 141 eV Kinetic Energy.

The non-magnetic system Ce [Figure 4, Ref 5-7].




Figure 4  
 Spin resolved results for a polycrystalline film of Ce deposited upon W(110). The photon energy was 127 eV. Top panel: spin polarized spectra. Middle panel: polarization. Lowest panel: Instrumental asymmetry. Circularly polarized photons were used to generate the spin dependence.




Figure 5

Spin-resolved photoemission spectra obtained with unpolarized He I( $h\nu = 21.2$  eV) light for normal emission from the valence bands of Pt(001). Upper panel for  $P_x$  component: The spin integrated total intensity  $I$  (black) and spin separated partial intensities  $I_+$  in red and  $I_-$  in blue are shown for 5x1 surface and destroyed surface. Since the measured spin polarization  $P_x$  is zero, the partial intensities  $I_+$  and  $I_-$  are identical. Lower panel for  $P_y$  component: The spin integrated total intensity  $I$  (black) and spin separated partial intensities  $I_+$  in red and  $I_-$  in blue are shown for 5x1 surface and destroyed surface. The measured spin polarizations  $P_y$  for 5x1 surface and destroyed surface are shown. Note the loss of the 5x1 reconstructed surface state intensity at the binding energy of 0.66 eV with the destruction of the surface order for both X and Y components.

Various recent publications have addressed the ongoing question of the nature of the Pu electronic structure. [9,10] One possibility is the existence of a Kondo-like shielding of the 5f electrons. [1] We propose that spin resolved PES is the most promising approach to resolving this question. [7,11]

#### Acknowledgements

Lawrence Livermore National Laboratory is operated by Lawrence Livermore National Security, LLC, for the U.S. Department of Energy, National Nuclear Security Administration under Contract DE-AC52-07NA27344. Work that was performed by UMR personnel was supported in part by the Office of Basic Energy Science at the U.S Department of Energy. Work that was performed by LLNL personnel was supported in part by the Office of Basic Energy Science at the U.S Department of Energy and Campaign 2 of WCI at LLNL.

- Present Address, Spring8 Synchrotron Radiation Facility, Japan

#### References

1. J.H. Shim, K. Haule and G. Kotliar, *Nature* **446**, 513 (2007).
2. J.G. Tobin, S.A. Morton, S.W. Yu, G.D. Waddill, I.K. Schuller, and S.A. Chambers, *J. Phys. Condens. Matter* **19**, 315218 (2007).
3. T. Komesu, G.D. Waddill, S.-W. Yu, M. Butterfield, and J.G. Tobin, "Spin-Orbit Effects in Spin-Resolved  $L_{2,3}$  Core Level Photoemission of 3d Ferromagnetic Thin Films," under preparation; J.G. Tobin and F.O. Schumann, *Surface Science* **478**, 211 (2001).
4. J.G. Tobin, S.A. Morton, S.W. Yu, T. Komesu, G.D. Waddill and P. Boyd, *Nuclear Instrum. Methods A* **582**, 165 (2007); J. D. W. Thompson, J. R. Neal, T. H. Shen, S. A. Morton, J. G. Tobin, G. D. Waddill, J.A.D. Matthew, D. Greig, and M. Hopkinson, submitted to PRB, 2007.
5. S.W. Yu et al, *Phys. Rev. B* **73**, 075116 (2006).
6. J.G. Tobin, S.A. Morton, B.W. Chung, S.W. Yu and G.D. Waddill, *Physica B* **378-380**, 925 (2006).
7. J.G. Tobin, S.W. Yu, T. Komesu, B.W. Chung, S.A. Morton, and G.D. Waddill, *Europhysics Lett.*, **77**, 17004 (2007).
8. S.-W. Yu and J.G. Tobin, *Surface Science Letters* **601**, L127 (2007).
9. J.G. Tobin et al, *Phys. Rev. B* **68**, 155109 (2003).
10. J.G. Tobin et al, *Phys. Rev. B* **72**, 085109 (2005).
11. J.G. Tobin, *J. Alloys Compds* **444-445**, 154 (2007).